

Supplementary data for article:

Baranac-Stojanović, M. Theoretical Analysis of the Rotational Barrier in Ethane: Cause and Consequences. *Structural Chemistry* **2015**, 26 (4), 989–996.

<https://doi.org/10.1007/s11224-014-0557-5>

Theoretical Analysis of the Rotational Barrier in Ethane: Cause and Consequences

Marija Baranac-Stojanović*

Faculty of Chemistry, University of Belgrade, Studentski trg 12-16, P.O.Box 158, 11000 Belgrade, Serbia

e-mail: mbaranac@chem.bg.ac.rs

Supplementary Material

Table of Contents

Table S1. Optimized C–C and C–H bond lengths, CCH and HCH bond angles for staggered and eclipsed conformations of ethane and differences between them.....	S2
Table S2. LMOEDA results for rigid rotation of staggered conformer.....	S3
Table S3. LMOEDA results for rigid rotation of eclipsed conformation.....	S3
Table S4. LMOEDA results for relaxed rotation of staggered conformer.....	S4

Table S1. Optimized C–C and C–H bond lengths, d_{CC} and d_{CH} (Å), CCH and HCH bond angles, τ_{CCH} and τ_{HCH} (°), for staggered (S) and eclipsed (E) conformations of ethane and differences between E and S (Δ_{E-S}).

Conformation	Method	d_{CC}	d_{CH}	τ_{CCH}	τ_{HCH}
S	HF/6-31G*	1.52736	1.08559	111.21533	107.67221
	HF/6-311+G**	1.52701	1.08631	111.19880	107.68976
	B3LYP/6-31G*	1.53066	1.09626	111.35280	107.52588
	B3LYP/6-311+G**	1.53089	1.09384	111.35547	107.52303
	MP2/6-311+G**	1.52925	1.09349	111.14651	107.74523
E	HF/6-31G*	1.54111	1.08460	111.65293	107.20411
	HF/6-311+G**	1.54104	1.08524	111.63206	107.22662
	B3LYP/6-31G*	1.54493	1.09515	111.78699	107.05943
	B3LYP/6-311+G**	1.54463	1.09271	111.77888	107.06822
	MP2/6-311+G**	1.54240	1.09264	111.62415	107.23510
Δ_{E-S}	HF/6-31G*	0.01375	-0.00099	0.43760	-0.46810
	HF/6-311+G**	0.01403	-0.00107	0.43326	-0.46314
	B3LYP/6-31G*	0.01427	-0.00111	0.43419	-0.46645
	B3LYP/6-311+G**	0.01374	-0.00113	0.42341	-0.45481
	MP2/6-311+G**	0.01315	-0.00085	0.47764	-0.51013

Table S2. LMOEDA results for rigid rotation of staggered conformer (B3LYP/6-311+G**).^a Values are in kcal/mol.

Torsional angle (°)	ΔE_{es}	$\Delta E_{\text{ex+rep}}$	ΔE_{pol}	ΔE_{disp}	ΔE_{int}	ΔE_{prep}	ΔE
0	-135.65	213.46	-160.93	-23.89	-107.01	18.49	-88.52
10	-135.64	213.28	-160.97	-23.89	-107.20	18.49	-88.71
20	-135.60	212.81	-161.07	-23.87	-107.72	18.49	-89.23
30	-135.55	212.15	-161.20	-23.84	-108.43	18.49	-89.94
40	-135.49	211.50	-161.34	-23.81	-109.14	18.49	-90.65
50	-135.46	211.02	-161.43	-23.79	-109.66	18.49	-91.17
60	-135.45	210.84	-161.47	-23.79	-109.85	18.49	-91.36

^a ΔE_{es} = electrostatic energy, ΔE_{ex} = exchange repulsion energy, ΔE_{pol} = polarization energy, ΔE_{disp} = dispersion energy, ΔE_{int} = interaction energy, ΔE_{prep} = preparation energy, ΔE = total bonding energy.

Table S3. LMOEDA results for rigid rotation of eclipsed conformation (B3LYP/6-311+G**).^a Values are in kcal/mol.

Torsional angle (°)	ΔE_{es}	$\Delta E_{\text{ex+rep}}$	ΔE_{pol}	ΔE_{disp}	ΔE_{int}	ΔE_{prep}	ΔE
0	-130.74	203.08	-156.86	-23.66	-108.19	19.53	-88.66
10	-130.73	202.92	-156.89	-23.65	-108.36	19.53	-88.83
20	-130.70	202.49	-156.99	-23.63	-108.83	19.53	-89.30
30	-130.65	201.89	-157.11	-23.61	-109.48	19.53	-89.95
40	-130.61	201.30	-157.24	-23.58	-110.13	19.53	-90.60
50	-130.58	200.85	-157.33	-23.57	-110.61	19.53	-91.08
60	-130.56	200.69	-157.36	-23.56	-110.79	19.53	-91.26

^a Labeling is the same as in Table S2.

Table S4. LMOEDA results for relaxed rotation of staggered conformer (B3LYP/6-311+G**).^a Values are in kcal/mol.

Torsional angle (°)	ΔE_{es}	$\Delta E_{\text{ex+rep}}$	ΔE_{pol}	ΔE_{disp}	ΔE_{int}	ΔE_{prep}	ΔE
0	-130.74	203.08	-156.86	-23.66	-108.19	19.53	-88.66
10	-130.93	203.27	-157.06	-23.67	-108.39	19.57	-88.82
20	-131.71	204.52	-157.83	-23.69	-108.70	19.39	-89.31
30	-132.87	206.46	-158.96	-23.72	-109.10	19.12	-89.98
40	-134.12	208.55	-160.17	-23.75	-109.48	18.82	-90.66
50	-135.11	210.26	-161.13	-23.78	-109.76	18.58	-91.18
60	-135.45	210.84	-161.47	-23.79	-109.85	18.49	-91.36

^a Labeling is the same as in Table S2.